

$\Lambda_b \rightarrow \Lambda + D(\bar{D}^0)$ decays and CP-violation

Fayyazuddin¹

Physics Department, Quaid-i-Azam University, Islamabad, Pakistan

Abstract

It is shown that interference of the amplitudes for the decays $\Lambda_b \rightarrow \Lambda D$ and $\Lambda_b \rightarrow \Lambda \bar{D}^0$ gives rise to CP -violation.

The decays $\Lambda_b \rightarrow \Lambda D^o$ and $\Lambda_b \rightarrow \Lambda \bar{D}^o$ are interesting for the following reasons. (i) The Hamiltonian for the decay $\Lambda_b \rightarrow \Lambda \bar{D}^o$ involves the weak phase $\hat{\gamma}$. This has direct implication for CP-Violation. (ii) The factorization contribution for these decays is suppressed due to the colour factor $C_2 + \xi C_1$ and the form factors $g_V(m_{D^2})$ and $g_A(m_{D^2})$. In reference [1], we have recently shown that $g_V(m_{D^2}) = g_A(m_{D^2}) = f_1(m_{D^2}) = -0.120$ and that the contribution of second form factor is negligible to the decay widths and asymmetry parameter α (In fact $f_2/f_1 = 0.129$). (iii) The baryon $\Lambda \Xi_c^o$ and Ξ_c^o poles contribute only to the decay $\Lambda_b \rightarrow \Lambda D^o$. The pole contribution can be evaluated in the W -exchange model [2].

The pole contributes to the p-wave amplitude and numerically its contribution is compareable to that of factorization. This has dramatic affect on the asymmetry parameter α . The asymmetry parameter $\alpha_{\bar{D}} = -1$, since only factorization contributes to this decay where as α_D can be as low as zero depending on the relative strength of the pole and factorization contributions.

In the Wolfenstein parameterization [3] of CKM matrix [4], the weak phases $\hat{\alpha}$, $\hat{\beta}$ and $\hat{\gamma}$ are given by $\tan \hat{\alpha} = \frac{\eta}{\eta^2 - \rho(1 - \rho)}$, $\tan \hat{\beta} = \frac{\eta}{1 - \rho}$ and $\tan \hat{\gamma} = \frac{\eta}{\rho}$. The weak phases have been denoted with $\hat{}$ to distinguish them from asymmetry parameters α , β and γ . In the two body non-leptonic decays of baryons, the angles ϕ and Δ defined as $\tan \phi = \beta/\gamma$ and $\tan \Delta = -\beta/\alpha$ [5] give direct information about the CP violation. Due to interference of amplitudes for the decays ΛD^o and $\Lambda \bar{D}^o$, the decay amplitudes for $\Lambda_b \rightarrow \Lambda D_{\pm}^o$, where $D_{\pm}^o = \frac{1}{\sqrt{2}}(D^o \pm \bar{D}^o)$ would make angles $\phi_{\pm}$ and

¹E-mail: fayyaz@hep-qau.sdnpk.undp.org

$\Delta_{\pm}$ non zero in the absence of final state interactions, if the pole contribution is not negligible. These decays have also been discussed in reference [6], but the overlap is minimal.

The decays $\Lambda_b \rightarrow \Lambda D^o$ and $\Lambda_b \rightarrow \bar{D}^o$ are described by the effective Hamiltonian [7]

$$\begin{aligned} H_{eff}(\Delta B = 1) &= \frac{G_F}{\sqrt{2}} [V_{cb}V_{us}^* (C_1 O_1^c + C_2 O_2^c) \\ &+ V_{ub}V_{cs}^* (C_1 O_1^u + C_2 O_2^u)], \end{aligned} \quad (1)$$

where C_i are Wilson coefficients evaluated at the normalization scale μ ; the current-current operators $O_{1,2}$ are

$$\begin{aligned} O_1^c &= (\bar{c}^\alpha b_\alpha)_{V-A} (\bar{s}^\beta u_\beta)_{V-A} \\ O_2^c &= (\bar{c}^\alpha b_\beta)_{V-A} (\bar{s}^\beta u_\alpha)_{V-A} \end{aligned} \quad (2)$$

and O_i^u are obtained through replacing c by u . Here α and β are $SU(3)$ color indices while $(\bar{c}^\alpha b_\alpha)_{V-A} = \bar{c}^\alpha \gamma_\mu (1 + \gamma_5) b_\alpha$ etc. We take [8]

$$\begin{aligned} C_1 &= 1.117 \\ C_2 &= -0.257. \end{aligned} \quad (3)$$

The factorization contributions to the s -wave and p -wave amplitudes A and B are given by

$$\begin{aligned} A_D &= -\frac{G_F}{\sqrt{2}} V_{cb}V_{us}^* (C_2 + \xi C_1) F_D (m_{\Lambda_b} - m_\Lambda) g_V(m_D^2) \\ &= \frac{G_F}{\sqrt{2}} (A\lambda^3) a_f \\ B_D &= \frac{G_F}{\sqrt{2}} (A\lambda^3) [b_f + b_p] \\ A_{\bar{D}} &= \frac{G_F}{\sqrt{2}} (A\lambda^3) (\rho - i\eta) a_f \\ B_{\bar{D}} &= \frac{G_F}{\sqrt{2}} (A\lambda^3) (\rho - i\eta) b_f \end{aligned} \quad (4)$$

where

$$\begin{aligned} a_f &= -(C_2 + \xi C_1) F_D (m_{\Lambda_b} - m_{\Lambda}) g_V \\ b_f &= (C_2 + \xi C_1) F_D (m_{\Lambda_b} + m_{\Lambda}) g_A \end{aligned} \quad (5)$$

In Eq. (3), we have also included the pole contribution b_p which is given by

$$\begin{aligned} B_p &= \frac{G_F}{\sqrt{2}} (A\lambda^3) b_p \\ &= - \left[\frac{g' \langle \Xi_c'^0 | H_W^{pc} | \Lambda_b \rangle}{m_{\Lambda_b} - m_{\Xi_c'^0}} + \frac{g \langle \Xi_c^0 | H_W^{pc} | \Lambda_b \rangle}{m_{\Lambda_b} - m_{\Xi_c^0}} \right] \end{aligned} \quad (6)$$

where $g' \equiv g_{\Xi_c'^0 \Lambda D}$ and $g \equiv g_{\Xi_c^0 \Lambda D}$ are strong coupling constants. In the W -exchange model [2], b_p is given by [9]

$$\begin{aligned} b_p &= \left[\frac{-\sqrt{3}g'}{m_{\Lambda_b} - m_{\Xi_c'^0}} + \frac{g}{m_{\Lambda_b} - m_{\Xi_c^0}} \right] d' \\ d' &= 5 \times 10^{-3} \text{GeV}^3 \end{aligned} \quad (7)$$

Note the important fact that $\Xi_c'^0$ and Ξ_c^0 poles contribute to the decay $\Lambda_b \rightarrow \Lambda D^0$ only. This is crucial in discussing the CP violation.

Let us symbolically write for the two channels of the decay:

$$|\Lambda_b\rangle = R_D |\Lambda D^0\rangle + R_{\bar{D}} |\Lambda \bar{D}^0\rangle \quad (8)$$

where, $R_D = A_D$ or B_D and $R_{\bar{D}} = A_{\bar{D}}$ or $B_{\bar{D}}$ for s or p -wave respectively. From Eq. (8), it is clear that the decay amplitudes for the states

$$\begin{aligned} D_+^0 &= \frac{1}{\sqrt{2}} (D^0 + \bar{D}^0), \\ D_-^0 &= \frac{1}{\sqrt{2}} (D^0 - \bar{D}^0), \end{aligned} \quad (9)$$

which are eigenstates of CP are given by

$$\begin{aligned} A_{\pm} &= \frac{1}{\sqrt{2}} \left[\frac{G_F}{\sqrt{2}} A\lambda^3 \right] [(1 \pm \rho) \mp i\eta] a_f \\ B_{\pm} &= \frac{1}{\sqrt{2}} \left[\frac{G_F}{\sqrt{2}} A\lambda^3 \right] [((1 \pm \rho) \mp i\eta) b_f + b_p] \end{aligned} \quad (10)$$

It is the presence of the pole contribution in the interference of the amplitudes that make $ImA_{\pm}^*B_{\pm}$ non zero, giving rise to CP violation. Now $ImA_{\pm}^*B_{\pm}$ is proportional to $\pm\eta(b_p/b_f)$. If b_p/b_f is not small i.e., if b_p is comparable with b_f , then $\beta_{\pm}$ may be experimentally measurable.

To fit the experimental branching ratio for the decay $\Lambda_b \rightarrow \Lambda J/\psi$, it was shown in reference [1], that with C_1 and C_2 as given in Eq. (3), the parameter $\xi = 1/N_c$ in $C_1 + \xi C_2$ must lie within the following limits:

$$0 \leq \xi \leq 0.125 \text{ or } 0.35 \leq \xi \leq 0.45 \quad (11)$$

These limits are consistent with the limit $0 \leq \xi \leq 0.5$ suggested by the combined analysis of the present CLEO data on $B \rightarrow h_1 h_2$ decay [8]. To give a crude estimate for the strong coupling constants g and g' , $SU(4)$ is used as a guide. In $SU(4)$

$$\begin{aligned} g' &= \sqrt{3/2}(g_f - g_d) = \sqrt{3/2}(1 - 2f)g_{\pi NN} \\ g &= -\sqrt{1/2}(g_f + g_d/3) = -\frac{1}{3\sqrt{2}}(1 + 2f)g_{\pi NN} \\ g'/g &= \sqrt{3}\frac{g_f - g_d}{g_f + g_d/3} = \sqrt{3}\frac{F - D}{F + D/3} \end{aligned} \quad (12)$$

Using $F = 0.446, D = 0.815$, we obtain

$$g'/g = -0.890, g = -0.493g_{\pi\pi N} \quad (13)$$

Now b_p/b_f can be easily obtained from Eqs. (5) and (7). For $\xi = 0, C_2 = -0.257, F_D = 0.200$ GeV, $g_A(m_D^2) = -0.120, m_{\Lambda_b} = 5.641$ GeV, $m_{\Lambda} = 1.116$ GeV, $m_{\Xi_c'^0} = 2.580$ GeV, $m_{\Xi_c^0} = 2.468$ GeV, $d' = 5 \times 10^{-3}$ GeV³, $g_{\pi NN} = 13.26$ and Eq. (13), we get

$$x \equiv b_p/b_f = -0.64 \quad (14)$$

For $\xi = 0.125$, one gets $x = -1.41$. Due to uncertainty in the parameter ξ and in the values of g' and g , we treat x as a free parameter in the range $-1 \leq x \leq 1$.

Using Eqns (4), (5), (7) and (10) and the experimental values of the masses given above, we obtain

$$\frac{\Gamma(\Lambda_b \rightarrow \Lambda D^0)}{\Gamma(\Lambda_b \rightarrow \Lambda \bar{D}^0)} = \frac{D}{\rho^2 + \eta^2} \quad (15)$$

$$\begin{aligned}\alpha_D &= \frac{-(1+x)}{D} \\ \alpha_{\bar{D}} &= -1\end{aligned}\tag{16}$$

$$D = 1 + 0.946x^2 + 0.473x^2\tag{17}$$

$$\tan \Delta_{\pm} = \frac{-[r_{\pm} \sin(\delta_p - \delta_s) + x \sin(\delta_p - \delta_s \pm \hat{\phi}, \hat{\beta})]}{[r_{\pm} \cos(\delta_p - \delta_s) + x \cos(\delta_p - \delta_s \pm \hat{\phi}, \hat{\beta})]}\tag{18}$$

$$\tan \phi_{\pm} = \frac{-[r_{\pm} \sin(\delta_p - \delta_s) + x \sin(\delta_p - \delta_s \pm \hat{\phi}, \hat{\beta})]}{[0.053r_{\pm} - 0.940x \cos(\hat{\phi}, \hat{\beta}) + 0.470\frac{x^2}{r_{\pm}}]}\tag{19}$$

where

$$\begin{aligned}r_{\pm} &= \sqrt{(1 \pm \rho)^2 + \eta^2} \\ \tan \hat{\phi} &= \frac{\eta}{1 + \rho}, \\ \tan \hat{\beta} &= \frac{\eta}{1 - \rho}\end{aligned}\tag{20}$$

Eqs. (15-20) are main results. Using [1]

$$\frac{\Gamma(\Lambda_b \rightarrow \Lambda \bar{D}^o)}{\Gamma(\Lambda_b \rightarrow \Lambda J/\Psi)} = (2.8 \times 10^{-3})(\rho^2 + \eta^2)\tag{21}$$

We can express Eq. (15) as

$$\frac{\Gamma(\Lambda_b \rightarrow \Lambda \bar{D}^o)}{\Gamma(\Lambda_b \rightarrow \Lambda J/\Psi)} = (2.8 \times 10^{-3})D\tag{22}$$

The above branching ratio and the asymmetry parameter α_D for various values of x are given in Table 1.

From Table 1 one can see that α_D can be as low as zero where as $\alpha_{\bar{D}} = -1$ independent of x .

In the absence of final state interaction i.e. $\delta_p - \delta_s = 0$, the angles $\Delta_{\pm}$ and $\phi_{\pm}$ as function of the parameter x given in Eqs. (18) and (19) are plotted in Figs. (1) and (2) respectively for the preferred values of (ρ, η) of reference [10] viz $(\rho, \eta) = (0.05, 0.36)$. Of course similar curves can be obtained for different values of (ρ, η) allowed by existing data.

Figs. (1) and (2) for the angles $\Delta_{\pm}$ and $\phi_{\pm}$ are a unique features of interference of the amplitudes for $\Lambda_b \rightarrow \Lambda D^o$ and $\Lambda \bar{D}^o$. The experimental values of α_D

would fix the value of the parameter x . Once x is known then Eqs. (18) and (19) [if $\delta_p - \delta_s \sim 0$] can be plotted as function of (ρ, η) . This may give some interesting information about CP violation.

However we have estimated $x = -0.64$ or -1.41 for $\xi = 0$ or $\xi = 0.125$. For these values of x and $(\rho, \eta) = (0.05, 0.36)$, we get

$$x = -0.64, \alpha_D = -0.61, \Delta_+ = 24^\circ, \Delta_- = -33^\circ, \phi_+ = 26^\circ, \phi_- = -28^\circ$$

$$x = -1.41, \alpha_D = 0.68, \Delta_+ = -64^\circ, \Delta_- = 59^\circ, \phi_+ = 48^\circ, \phi_- = -53^\circ$$

It is clear that determination of α_D in future experiment will be by itself interesting even though the experimental observation of CP violation in these decays may be difficult in near future.

I am grateful to Riazuddin for discussions.

x	α_D	$\left[\frac{\Gamma(\Lambda_b \rightarrow \Lambda \bar{D}^0)}{\Gamma(\Lambda_b \rightarrow \Lambda J/\Psi)} \right] \times 10^3$
-1	0	1.5
-0.5	- 0.77	1.8
0	-1.0	2.8
0.5	- 0.94	4.4
1	- 0.83	6.8

Table - I **The decay parameter α_D and the braching ratio $\frac{\Gamma(\Lambda_b \rightarrow \Lambda \bar{D}^0)}{\Gamma(\Lambda_b \rightarrow \Lambda J/\Psi)}$.**




Figure 1: The angle $\Delta_{\pm} = \arctan(-\beta_{\pm}/\alpha_{\pm})$ as a function of x [See Eq. (18), $(\delta_p - \delta_s) = 0$] for $(\rho, \eta) = (0.05, 0.36)$. Solid line gives Δ_+ and dotted line gives Δ_- .




Figure 2: The angle $\phi_{\pm} = \arctan(\beta_{\pm}/\gamma_{\pm})$ as a function of x [See Eq. (19), $(\delta_p - \delta_s) = 0$] $(\rho, \eta) = (0.05, 0.36)$. Solid line gives ϕ_+ and dotted line gives ϕ_- .

References

- [1] Fayyazuddin and Riazuddin, " Two body non-leptonic Λ_b decays in quark model with factorization ansatz" in press *Phys. Rev. D* (1998).
- [2] Riazuddin and Fayyazuddin, *Phys. Rev. D* **18** (1978) 1578; **19**(1978)1630E
- [3] L. Wolfenstein, *Phys. Rev. Lett* **51**(1983)1945
- [4] N. Cabbibo, *Phys. Rev. Lett.* **10**(1963)531; M. Kobayashi and K. Maskawa, *Prog. Theor. Phys.* **49**(1973)652
- [5] Particle Data Group, Review of Particle properties, *Phys. Rev. D* **54**, Part-1 (1996)
- [6] I. Dunietz, *Z. Phys' C* **56**(1992)129
- [7] For a review, see G. Buchalla, A.J. Buras and M.E. Lauternbacher, *Rev. Mod. Phys.* **68**(1996)1125
- [8] A. Ali and C. Greub, DESY Preprint, DESY 97-126 [hep-ph/9707251]
- [9] Fayyazuddin and Riazuddin, *Phys. Rev. D* **55** (1997)255
- [10] A. Ali and D. London, DESY Preprint, DESY 96-140 [hep-ph/9607392]